



Case Report

Pancreatic Rest Complicated by Actinomyces Gastric Abscess in a Young Male: A Case ReportAyah Obeid^{1,*}, Frank Lin¹, Anuraag Bandi¹, Loveleen Sidhu², Sinan Kutty², Gurshawn Singh², Lisa Stoll³

1-Internal Medicine Department, St. Luke's University Health Network, Bethlehem, PA, USA

2-Gastroenterology Department, St. Luke's University Health Network, Bethlehem, PA, USA

3-Pathology Department, St. Luke's University Health Network, Bethlehem, PA, USA

ARTICLE INFO

Article history:

Received 25 Apr. 2025

Received in revised form 13 May. 2025

Accepted 22 May. 2025

Published 15 Jun. 2023

Keywords:

Endoscopic ultrasound (EUS)

Gastric abscess

Heterotopic pancreas

Pancreatic rest

ABSTRACT

Pancreatic rest, or ectopic pancreatic tissue, is a rare condition. It is characterized by pancreatic tissue outside its usual location, most commonly in the gastric antrum or proximal small intestine. We present a rare case of a 21-year-old male with recurrent epigastric pain, vomiting, and fever. Imaging and endoscopic ultrasound (EUS) identified a subepithelial lesion with features consistent with pancreatic rest. Subsequent fine-needle aspiration (FNA) grew *Actinomyces* and *Streptococcus*. Given persistent symptoms and incomplete resolution despite prolonged antibiotics, he underwent partial gastrectomy. This case highlights an unusual infectious complication of pancreatic rest with *Actinomyces* and underscores the need to consider surgical intervention in cases refractory to medical therapy.

1. Introduction

Pancreatic rest is a condition where pancreatic tissue is located outside its usual anatomical site. It is also called ectopic, heterotopic, or aberrant pancreatic tissue. This tissue lacks an anatomical or vascular connection to the main pancreas yet retains its own separate blood and nerve supply [1, 2]. This condition predominantly affects the gastrointestinal tract, particularly the gastric antrum and proximal small intestine, with an incidence ranging from 0.55% to 13% in autopsy series [1]. While pancreatic rests are often asymptomatic and discovered incidentally during endoscopic or imaging studies, common symptoms include nausea, vomiting, epigastric pain, weight loss, gastrointestinal bleeding, and gastric outlet obstruction. Rarely, it can lead to complications such as gastric ulceration, pancreatitis, obstructive jaundice when near the Ampulla of Vater, or even acute perforation due to inflammation in the stomach or duodenum [2, 3]. Gastric abscesses secondary to pancreatic rest are particularly rare, with few reported cases describing this phenomenon [4, 5]. These abscesses are believed to result from localized inflammation or necrosis of ectopic acinar tissue. Identifying pancreatic rest as the source of a gastric abscess is clinically challenging, particularly when the initial presentation mimics more common gastric subepithelial lesions such as gastrointestinal stromal tumors (GISTs) or duplication cysts [2, 6]. Malignant transformation is uncommon, occurring in 0.7% to 1.8% of cases [3]. For example, a case report highlighted a pancreatic

rest presenting as a subepithelial nodule, with histological analysis revealing pancreatic intraepithelial neoplasia [7]. We report a rare case of a young male with recurrent gastric abscess formation due to pancreatic rest, complicated by *Actinomyces* and alpha-hemolytic *Streptococcus* infection. To our knowledge, this is among the few reported cases of infected pancreatic rest and the first with hematogenous *Actinomyces* seeding from dental work.

2. Case Presentation

A 21-year-old male with no significant past medical history initially presented to the emergency department (ED) with epigastric abdominal pain radiating to the back, early satiety, and vomiting. Laboratory workup showed white blood cells (WBC) 15×10^3 , lipase 26 U/L, and alanine aminotransferase 16 U/L. A computed tomography (CT) scan of the abdomen and pelvis was done, and it showed mild edematous and inflammatory stranding adjacent to the distal stomach. He was discharged with a prescription for omeprazole and metoclopramide and advised to follow up with gastroenterology. One week later, he returned with worsening symptoms and a fever (101°F). Repeat labs showed leukocytosis 18×10^3 , Lipase 26 U/L, an alkaline phosphatase of 76 U/L, aspartate aminotransferase of 23 U/L, alanine aminotransferase of 29 U/L, and lipase of 26 U/L. A repeat CT scan with contrast demonstrated mass-like thickening of the gastric antral wall with luminal compression and adjacent stranding, suggesting an inflammatory phlegmon with no drainable collection (**Figure 1**). He was admitted and started on ceftriaxone and Metronidazole. Esophagogastroduodenoscopy (EGD) revealed large subepithelial extrinsic compression in the antrum, with the suspected fluid collection, possibly a walled-off abscess (**Figure 2**). No ulceration was observed, and biopsies were not obtained due to the soft consistency of the lesion. He was discharged on Metronidazole and Cefdinir for 7 days with plans for a repeat CT scan and EGD in 6 weeks. A follow-up CT scan

*Corresponding author: Ayah Obeid, Internal Medicine Department, St. Luke's University Health Network, Bethlehem, PA, USA. Email: obeidayah@yahoo.com

Published in collaboration with the American Society for Inclusion, Diversity, and Equity in Healthcare (ASIDE). ISSN (Print): 3066-4004, ISSN (Online): 3066-4012 – See front matter © 2025 ASIDE Gastroenterology. This work is licensed under a Creative Commons Attribution 4.0 International License. Hosted by ASIDE Journals.

Citation: Obeid A, Lin F, Bandi A, et al. Pancreatic Rest Complicated by Actinomyces Gastric Abscess in a Young Male: A Case Report. ASIDE GI. 1(3):1-4. doi:10.71079/ASIDE.GI.06012584

showed interim resolution of the antral mass. The EGD showed a 15 mm submucosal nodule along the greater curvature of the antrum concerning pancreatic rest. EUS revealed an extended area of heterogenous echotexture extending from the muscularis propria to the submucosa, consistent with pancreatic rest (**Figure 3**). Five months later, the patient presented to the ED with recurrent epigastric pain, nausea, and diarrhea. Labs showed WBC $15 \times 10^3/\mu\text{L}$, lipase 26 U/L, and amylase 43 U/L. An abdominal-pelvic CT scan with contrast showed a 2.4 cm fluid-containing structure in the anterior/caudal wall of the gastric antrum near the pylorus, suspicious of an abscess versus a walled-off ulcer (**Figure 4**). Ceftriaxone and metronidazole were reinitiated. A repeat EUS showed an irregular, oval, and anechoic cyst measuring 27 mm x 21 mm, with debris present and well-defined, smooth margins (**Figure 5**). Fine-needle aspiration revealed no malignant cells, with abundant neutrophils and histiocytes. Culture grew alpha-hemolytic *Streptococcus* and *Actinomyces* species. He was discharged on a 7-day course of cefpodoxime and metronidazole. A repeat CT scan four weeks later showed reduced size of the gastric wall fluid collection, with no new findings. Due to persistence, he was referred to Infectious Diseases and started on cefdinir and doxycycline. Imaging showed a 2.2×1.9 cm gastric antral mass without fluid two months later. Cefdinir was stopped, and doxycycline was continued for six months to treat *Actinomyces*. He later reported a recent root canal prior to symptom onset, suggesting an odontogenic source. Given the persistence of the lesion despite multiple courses of antibiotics and concerns over prolonged antibiotic use in a young patient, he underwent partial gastrectomy. Preoperative CT showed a persistent submucosal mass measuring $2.3 \times 2.0 \times 2.0$ cm. A surgical biopsy showed a submucosal abscess involving a small bowel and stomach with foreign-body giant cells and clear margins (**Figure 6**). At the one-year follow-up, the patient remained asymptomatic and denied any gastrointestinal symptoms, and no follow-up imaging was performed due to the absence of clinical concerns.




Figure 1: Mass-like thickening of the wall of the gastric antrum with luminal compression and adjacent stranding.

3. Discussion

Pancreatic rest is typically found incidentally during endoscopic imaging and is considered a gastric subepithelial lesion when found incidentally in the stomach. The differential for gastric subepithelial lesions can be broad, but it is important to consider pancreatic rest. Pancreatic rest can be found throughout the gastrointestinal tract; however, it is more commonly found in the stomach, particularly in the distal stomach along the greater curvature of the antrum [7]. Our patient's endoscopic ultrasound revealed a submucosal nodule with heterogenous echotexture extending from the muscularis propria to the submucosa of the greater curvature of the antrum of the stomach consistent with pancreatic rest. The classic




Figure 2: Large subepithelial extrinsic compression towards the antrum. Suspected fluid collection, possibly a walled-off abscess on the other side of this. No clear ulceration was seen.




Figure 3: A: Antral nodule with redundant folds. General appearance consistent with pancreatic rest. B: Area of thickened fold. The muscularis propria thickened to 12 mm with an extended area of heterogenous echotexture extending from the muscularis propria to the submucosa, consistent with pancreatic rest.



Figure 4: Rim-enhancing 2.4 cm fluid-containing structure is observed, including the anterior/caudal wall of the antrum of the stomach near the pylorus.

appearance of pancreatic rest on endoscopic ultrasound is described as hypoechoic or mixed echogenicity with heterogeneity within the second, third, or fourth layer (muscularis mucosa, submucosa, and muscularis propria, respectively) [4]. The heterogenous appearance is attributed to the presence of acini within the ectopic pancreatic tissue [7]. EUS is the preferred imaging study of choice to distinguish pancreatic rest from other gastric sub-mucosal lesions, with GISTs and leiomyomas being the most common [5]. The finding within the stomach can commonly be mistaken for malignancy; however, malignant features of an intramural lesion seen on EUS are typically greater than 4 cm in size, echogenic foci greater than 3 mm, cystic areas within the lesion, irregular borders, and the




Figure 5: Irregular, oval, and anechoic cyst measuring 27 mm x 21 mm with debris present, well-defined margins, and smooth margins.




Figure 6: A: HE, 10X: Brunner glands of the duodenum (upper right) with adjacent acute and chronic inflammation, including multinucleated giant cells consistent with abscess. B: HE, 20X; Abundant neutrophils with scattered multinucleated giant cells (arrow).

presence of adjacent lymph nodes with malignant pattern [6]. Our patient did not have any of these malignant features. Although pancreatic rest is typically asymptomatic, it can result in complications, including acute or chronic pancreatitis, pancreatic necrosis, pseudocyst, abscess, gastric outlet obstruction, and carcinoma [5, 8]. Gastric abscess formation secondary to pancreatic rest has been reported only in a few cases. Alastal et al. reported a patient with an ectopic pancreas complicated by recurrent abscess and pancreatitis, ultimately requiring surgical intervention [9]. Similarly, Berry et al. described a heterotopic pancreas initially mistaken for malignancy, necessitating resection [10]. However, neither case identified *Actinomyces* or a suspected odontogenic source. Our case is unique in that the abscess was polymicrobial, including *Actinomyces*, and may have originated from hematogenous spread following a recent root canal procedure, a mechanism not previously reported. Furthermore, while Alastal's case involved abscesses in the setting of ectopic pancreatitis [9], our patient lacked clinical or imaging features of pancreatitis. Instead, recurrent abscesses formed without enzymatic inflammation, possibly due to localized tissue necrosis and infection within the pancreatic rest itself. Histopathology supported the diagnosis, with foreign-body giant cells reflecting chronic inflammation indicative of a prolonged immune response. Their presence suggests the body's attempt to wall off or isolate the ectopic pancreatic tissue, further supporting the chronicity and infectious etiology of the lesion. These cells form granulomas that may occur in response to tissue injury in ectopic pancreatic rests [11]. Management of such cases typically begins with antibiotic therapy, as was initially attempted in our patient. However, despite multiple courses of antibiotics and temporary clinical improvement, imaging showed persistence of the lesion, ultimately necessitating partial gastrectomy. This case underscores the importance of recognizing rare infectious complications of

pancreatic rest. It demonstrates how multidisciplinary evaluation, including imaging, endoscopy, pathology, infectious disease, and surgical intervention, can lead to successful outcomes. It also contributes to a better understanding of *Actinomyces*-associated gastric abscesses, which are exceedingly rare and may present in atypical ways. This case report is limited by the lack of histological confirmation of pancreatic acini in the initial diagnosis, although imaging and clinical features were consistent. Additionally, follow-up beyond one year would be beneficial to assess for long-term recurrence.

4. Conclusions

This case illustrates a rare but clinically significant complication of pancreatic rest and highlights the need for timely recognition and coordinated care. Awareness of atypical infectious presentations can guide appropriate interventions and improve patient outcomes.

Conflicts of Interest

The authors declare that they have no competing interests that could have influenced the objectivity or outcome of this investigation.

Funding Source

None

Acknowledgements

None

Informed consent statement

Consent was obtained from the patient and family to publish this case report and images. All relevant information and confidentiality rights were explained, and identifying details have been anonymized.

Large Language Model Statement

None

Authors Contribution Statement

AO provided study leadership and led manuscript writing and editing; FL and AB contributed to manuscript writing; LoS supervised the project and reviewed the manuscript; SK and GS performed advanced scoping for the case and reviewed the manuscript and figures; LiS reviewed pathology slides and provided their descriptions. All authors reviewed and approved the final manuscript and ensured its accuracy and integrity.

Data Availability Statement

All information presented in this case report is included within the manuscript. If further details are required, please contact the corresponding author.

References

- Masoodi I, Al-Lehibi A, Almohaimeed K, Hussain S. Pancreatic Rest - An Unusual Cause of Dyspepsia: A Case Report with Literature

- Review. Saudi J Med Med Sci. 2016;4(3):225-8. Masoodi, Ibrahim Al-Lehibi, Abed Almohaimeed, Khalid Hussain, Shabnum eng Case Reports India 2016/09/01 Saudi J Med Med Sci. 2016 Sep-Dec;4(3):225-228. doi: 10.4103/1658-631X.188261. Epub 2016 Aug 11. [PMCID: PMC6298353, <https://doi.org/10.4103/1658-631X.188261>].
2. Kim DU, Lubner MG, Mellnick VM, Joshi G, Pickhardt PJ. Heterotopic pancreatic rests: imaging features, complications, and unifying concepts. *Abdom Radiol (NY)*. 2017;42(1):216-25. Kim, David U Lubner, Meghan G Mellnick, Vincent M Joshi, Gayatri Pickhardt, Perry J eng Review 2016/08/22 *Abdom Radiol (NY)*. 2017 Jan;42(1):216-225. doi: 10.1007/s00261-016-0874-9. [<https://doi.org/10.1007/s00261-016-0874-9>].
 3. Leung G, Mills J, Bucobo JC, Docimo S. Evaluation and management of a pancreatic rest noted during pre-bariatric surgery screening endoscopy. *Surg Endosc*. 2021;35(2):536-61. Leung, Galen Mills, John Bucobo, Juan Carlos Docimo, Salvatore eng Review Germany 2020/10/03 *Surg Endosc*. 2021 Feb;35(2):536-561. doi: 10.1007/s00464-020-08040-2. Epub 2020 Oct 1. [<https://doi.org/10.1007/s00464-020-08040-2>].
 4. Humphris JL, Jones DB. Subepithelial mass lesions in the upper gastrointestinal tract. *J Gastroenterol Hepatol*. 2008;23(4):556-66. Humphris, Jeremy L Jones, D Brian eng Review Australia 2007/12/19 *J Gastroenterol Hepatol*. 2008 Apr;23(4):556-66. doi: 10.1111/j.1440-1746.2007.05232.x. Epub 2007 Dec 14. [<https://doi.org/10.1111/j.1440-1746.2007.05232.x>].
 5. Iwahashi S, Nishi M, Yoshimoto T, Kashihara H, Takasu C, Tokunaga T, et al. A case of gastric heterotopic pancreas with gastroduodenal invagination. *Surg Case Rep*. 2019;5(1):110. Iwahashi, Shoko Nishi, Masaaki Yoshimoto, Toshiaki Kashihara, Hideya Takasu, Chie Tokunaga, Takuya Miyatani, Tomohiko Higashijima, Jun Yoshikawa, Kozo Wada, Yuma Bando, Yoshimi Shimada, Mitsuo eng Japan 2019/07/12 *Surg Case Rep*. 2019 Jul 10;5(1):110. doi: 10.1186/s40792-019-0669-7. [PMCID: PMC6620227, <https://doi.org/10.1186/s40792-019-0669-7>].
 6. Pih GY, Kim DH. Endoscopic Ultrasound-Guided Fine Needle Aspiration and Biopsy in Gastrointestinal Subepithelial Tumors. *Clin Endosc*. 2019;52(4):314-20. Pih, Gyu Young Kim, Do Hoon eng Korea (South) 2019/08/03 *Clin Endosc*. 2019 Jul;52(4):314-320. doi: 10.5946/ce.2019.100. Epub 2019 Jul 30. [PMCID: PMC6680013, <https://doi.org/10.5946/ce.2019.100>].
 7. Bahirwani J, Duarte-Chavez R, Stoll L, Matin A. Endoscopic Mucosal Resection of Pancreatic Rest Presenting as a Sub-epithelial Nodule in the Gastric Antrum. *Cureus*. 2023;15(12):e50713. Bahirwani, Janak Duarte-Chavez, Rodrigo Stoll, Lisa Matin, Ayaz eng Case Reports 2024/01/18 *Cureus*. 2023 Dec 18;15(12):e50713. doi: 10.7759/cureus.50713. eCollection 2023 Dec. [PMCID: PMC10792705, <https://doi.org/10.7759/cureus.50713>].
 8. Wawrzynski J, Leon L, Shah SA, Adrain A, Goldstein LJ, Feller E. Gastric Heterotopic Pancreas Presenting as Abdominal Pain with Acute and Chronic Pancreatitis in the Resected Specimen. *Case Rep Gastrointest Med*. 2019;2019(2021712):2021712. Wawrzynski, Joseph Leon, Lauren De Shah, Samir A Adrain, Alyn Goldstein, Lisa J Feller, Edward eng Case Reports 2019/04/09 *Case Rep Gastrointest Med*. 2019 Mar 6;2019:2021712. doi: 10.1155/2019/2021712. eCollection 2019. [PMCID: PMC6431401, <https://doi.org/10.1155/2019/2021712>].
 9. Alastal Y, Khalil B, Singh S, Almadani SB. Ectopic Pancreas in the Gastric Antrum Wall Complicated by Ectopic Pancreatitis and Persistent Gastric Abscess. *ACG Case Rep J*. 2018;5:e34. Alastal, Yaseen Khalil, Basmah Singh, Surendra Almadani, Sameh B eng Case Reports 2018/05/19 *ACG Case Rep J*. 2018 May 9;5:e34. doi: 10.14309/crj.2018.34. eCollection 2018. [PMCID: PMC5948316, <https://doi.org/10.14309/crj.2018.34>].
 10. Berry R, Rahal HK, Ho W. Pancreatic Heterotopia Found in the Gastric Antrum Mistaken for Malignancy. *ACG Case Rep J*. 2019;6(10):e00232. Berry, Rani Rahal, Harman K Ho, Wendy eng Case Reports 2019/12/14 *ACG Case Rep J*. 2019 Oct 14;6(10):e00232. doi: 10.14309/crj.0000000000000232. eCollection 2019 Oct. [PMCID: PMC6855548, <https://doi.org/10.14309/crj.0000000000000232>].
 11. Wang WG, Zhang Y, Wang L, Chen Y, Tian BL. Chronic pancreatic inflammatory granuloma caused by foreign body presenting as a pancreatic pseudotumor: A case report and literature review. *Pancreatol*. 2015;15(5):573-5. Wang, Wei-Guo Zhang, Yi Wang, Li Chen, Yang Tian, Bo-Le eng Case Reports Review Switzerland 2015/06/23 *Pancreatol*. 2015 Sep-Oct;15(5):573-575. doi: 10.1016/j.pan.2015.05.474. Epub 2015 Jun 5. [<https://doi.org/10.1016/j.pan.2015.05.474>].