

Contents

Supplemental Table 1 Angina Pectoris–related Deaths, Stratified by Sex and Race, in Older Adults in the United States, 1999 to 2023.	2
Supplemental Table 2 Angina Pectoris–related Mortality, Stratified by Place of Death in Older Adults in the United States, 1999 to 2023.....	3
Supplemental Table 3 Annual percent change (APC) of Angina Pectoris–related Age-Adjusted Mortality Rates per 100,000, to 2000 U.S. population in Older Adults in the United States, 1999 to 2023.	4
Supplemental Table 4: Overall and Sex - Stratified Angina Pectoris – Related Age-Adjusted Mortality Rates per 100,000, to 2000 U.S. population in Older Adults in the United States, 1999 to 2023.	7
Supplemental Table 5 Angina Pectoris–related Age-Adjusted Mortality Rates per 100,000, to 2000 U.S. population Stratified by Race in Older Adults in the United States, 1999 to 2023.	8
Supplemental Table 6 Angina Pectoris –related Crude Mortality Rates per 100,000, to 2000 U.S. population Stratified by Age Group in Adults in the United States, 1999 to 2023.	9
Supplemental Table 7 Angina Pectoris–related Age-Adjusted Mortality Rates per 100,000, to 2000 U.S. population Stratified by Census Region in Older Adults in the United States, 1999 to 2023.	10
Supplemental Table 8 Angina Pectoris–related Age-Adjusted Mortality Rates per 100,000, to 2000 U.S. population Stratified by Urban-Rural Classification in Older Adults in the United States, 1999 to 2020.	13

Supplemental Table 1 Angina Pectoris–related Deaths, Stratified by Sex and Race, in Older Adults in the United States, 1999 to 2023.

Year	Overall	Women	Men	NH White	NH Black or African American	Hispanic or Latino	NH Asian or Pacific Islander	NH American Indian or Alaska Native
1999	2,581	1,562	1,019	2,395	157	69	20	Missing
2000	2,564	1,557	1,007	2,363	175	67	23	Missing
2001	2,219	1,323	896	2,027	156	64	29	Missing
2002	1,950	1,195	755	1,765	152	53	27	Missing
2003	1,636	968	668	1,512	104	51	18	Missing
2004	1,367	795	572	1,243	98	53	25	Missing
2005	1,239	732	507	1,149	71	43	16	Missing
2006	1,048	607	441	967	64	34	13	Missing
2007	965	567	398	881	53	47	24	Missing
2008	899	513	386	819	65	51	13	Missing
2009	827	470	357	750	52	37	19	Missing
2010	816	466	350	730	57	48	21	Missing
2011	762	401	361	694	49	44	14	Missing
2012	811	441	370	726	67	47	11	Missing
2013	823	440	383	739	55	47	25	Missing
2014	798	412	386	721	60	76	14	Missing
2015	895	477	418	786	80	78	24	Missing
2016	1,278	657	621	1,156	82	115	33	Missing
2017	1,457	726	731	1,289	113	104	54	Missing
2018	1,590	821	769	1,422	113	138	39	16
2019	1,695	860	835	1,497	130	138	56	12
2020	2,050	1,003	1,047	1,806	159	222	69	16
2021	2,189	1,039	1,150	1,913	174	212	90	11
2022	2,108	1,024	1,084	1,876	145	198	75	10
2023	1,977	935	1,042	1,734	151	173	71	17
Total	36,544	19,991	16,553	32,960	2,582	2,209	823	82

NH, non-Hispanic.

Supplemental Table 2 Angina Pectoris–related Mortality, Stratified by Place of Death in Older Adults in the United States, 1999 to 2023.

Place of Death	Deaths
Medical Facility - Inpatient	11,130
Medical Facility - Outpatient or ER	2,672
Medical Facility - Dead on Arrival	222
Medical Facility - Status unknown	40
Decedent's home	11,269
Hospice facility	810
Nursing home/long-term care	8,843
Other	1,507
Place of death unknown	49
Total	36,542

Supplemental Table 3 Annual percent change (APC) of Angina Pectoris–related Age-Adjusted Mortality Rates per 100,000, to 2000 U.S. population in Older Adults in the United States, 1999 to 2023.

Year	APC (95% CI)
Overall	
1999-2009	−13.48 (−16.07 to −12.33)
2009-2014	−0.42 (−10.8 to 7.78)
2014-2021	13.14 (11.11 to 20.92)
2021-2023	−8.24 (−15.31 to 0.18)
Women	
1999-2001	−7.66 (−14.46 to −1.6)
2001-2006	−16.78 (−21.62 to −7.31)
2006-2013	−6.96 (−10.18 to 7.46)
2013-2021	11.81 (10.18 to 15.7)
2021-2023	−8.46 (−14.7 to −0.57)
Male	
1999-2009	−13.09 (−14.9 to −12.04)
2009-2014	0.79 (−7.9 to 7.44)
2014-2021	13.74 (11.73 to 19.38)
2021-2023	−7.3 (−13.19 to 1.04)
NH White	
1999-2008	−13.74 (−17.03 to −12.51)
2008-2013	−4.43 (−10.88 to 6.85)
2013-2021	12.54 (10.87 to 18.15)
2021-2023	−7.73 (−14.7 to 0.83)
NH Black or African American	
1999-2002	−5.2 (−11.26 to 6.91)
2002-2006	−21.96 (−29.53 to −16.64)
2006-2013	−3.99 (−11.1 to 5.37)
2013-2021	11.34 (8.89 to 23.93)
2021-2023	−8.1 (−18.7 to 3.84)
Hispanic or Latino	
1999-2011	−9.01 (−12.52 to −6.44)

2011-2021	12.81 (10.52 to 19.67)
2021-2023	−13.54 (−24.26 to 1.44)
Metropolitan	
1999-2008	−13.78 (−17.54 to −12.46)
2008-2013	−2.96 (−11.03 to 6.85)
2013-2020	12.38 (9.82 to 18.77)
Non-metropolitan	
1999-2001	−1.45 (−7.79 to 3.37)
2001-2007	−16.72 (−20.8 to −15.24)
2007-2014	−5.78 (−8.51 to −2.61)
2014-2020	13.52 (10.78 to 17.25)
Northeast	
1999-2009	−15.48 (−19.94 to −14.09)
2009-2014	−6.27 (−14.37 to 9.16)
2014-2020	16.6 (12.07 to 30.23)
2020-2023	−6.85 (−16.37 to 0.02)
Midwest	
1999-2001	−9.51 (−15.03 to −3.35)
2001-2009	−16.01 (−22.35 to −4.49)
2009-2014	−2.41 (−11.29 to 9.41)
2014-2021	13.08 (10.75 to 20.93)
2021-2023	−9.04 (−16.68 to 0.13)
South	
1999-2004	−14.14 (−17.89 to −12.49)
2004-2014	−0.97 (−13.05 to 8.28)
2014-2021	13.9 (10.96 to 24.55)
2021-2023	−4.31 (−13.39 to 6.59)
West	
1999-2011	−7.29 (−9.59 to −5.43)
2011-2021	10.42 (8.35 to 17.27)
2021-2023	−9.54 (−19 to 4.53)
Age 65-74 Years	
1999-2001	−5.2 (−10.38 to −0.67)
2001-2006	−17.21 (−20.73 to −15.67)
2006-2013	−4.99 (−7.44 to −2.43)

2013-2021	11.49 (10.4 to 13.2)
2021-2023	-11.02 (-14.84 to -4.08)
Age 75-84 Years	
1999-2001	-6.45 (-13.37 to 0.55)
2001-2006	-17.65 (-23.21 to -14.74)
2006-2013	-5.28 (-9.02 to 2.35)
2013-2021	12.38 (10.72 to 16.5)
2021-2023	-9.93 (-16.53 to -1.83)
Age 85+ Years	
1999-2009	-13.44 (-14.95 to -12.47)
2009-2014	-1.33 (-7.99 to 5.63)
2014-2021	13.9 (11.95 to 19.99)
2021-2023	-6.09 (-11.82 to 3.23)

APC = annual percent change; NH = non-Hispanic.

Supplemental Table 4: Overall and Sex-Stratified Angina Pectoris–Related Age-Adjusted Mortality Rates per 100,000, to 2000 U.S. population in Older Adults in the United States, 1999 to 2023.

Year	Overall	Women	Men
1999	7.52 (7.23-7.81)	6.95 (6.6-7.29)	8.22 (7.7-8.73)
2000	7.34 (7.05-7.62)	6.87 (6.53-7.22)	8.03 (7.53-8.54)
2001	6.27 (6.01-6.53)	5.83 (5.52-6.15)	6.95 (6.49-7.41)
2002	5.44 (5.2-5.68)	5.21 (4.91-5.5)	5.76 (5.34-6.17)
2003	4.5 (4.29-4.72)	4.14 (3.88-4.4)	5.05 (4.66-5.44)
2004	3.74 (3.54-3.93)	3.39 (3.15-3.63)	4.15 (3.81-4.5)
2005	3.34 (3.16-3.53)	3.04 (2.81-3.26)	3.64 (3.32-3.96)
2006	2.73 (2.57-2.9)	2.46 (2.26-2.65)	3.07 (2.78-3.35)
2007	2.45 (2.3-2.6)	2.26 (2.08-2.45)	2.72 (2.45-2.99)
2008	2.25 (2.11-2.4)	2.06 (1.88-2.24)	2.56 (2.3-2.82)
2009	2.02 (1.88-2.16)	1.88 (1.71-2.06)	2.28 (2.04-2.52)
2010	1.96 (1.82-2.09)	1.81 (1.65-1.98)	2.21 (1.98-2.44)
2011	1.8 (1.67-1.93)	1.53 (1.37-1.68)	2.16 (1.94-2.39)
2012	1.85 (1.72-1.98)	1.63 (1.48-1.79)	2.18 (1.95-2.4)
2013	1.84 (1.71-1.97)	1.58 (1.43-1.73)	2.21 (1.98-2.43)
2014	1.71 (1.59-1.83)	1.47 (1.32-1.61)	2.1 (1.88-2.31)
2015	1.91 (1.79-2.04)	1.67 (1.51-1.82)	2.21 (1.99-2.42)
2016	2.65 (2.5-2.79)	2.24 (2.06-2.41)	3.25 (2.99-3.5)
2017	2.93 (2.77-3.08)	2.42 (2.24-2.6)	3.65 (3.38-3.92)
2018	3.16 (3-3.31)	2.68 (2.49-2.86)	3.72 (3.46-3.99)
2019	3.22 (3.07-3.38)	2.73 (2.55-2.92)	3.94 (3.67-4.21)
2020	3.87 (3.7-4.04)	3.15 (2.96-3.35)	4.85 (4.55-5.14)
2021	4.38 (4.2-4.57)	3.56 (3.34-3.78)	5.53 (5.2-5.86)
2022	4 (3.83-4.17)	3.21 (3.01-3.41)	5.06 (4.76-5.37)
2023	3.73 (3.57-3.9)	3.02 (2.82-3.21)	4.75 (4.46-5.05)

Supplemental Table 5 Angina Pectoris–related Age-Adjusted Mortality Rates per 100,000, to 2000 U.S. population Stratified by Race in Older Adults in the United States, 1999 to 2023.

Year	NH White	NH Black or African American	Hispanic or Latino	NH Asian or Pacific Islander	NH American Indian or Alaska Native
1999	7.75 (7.44-8.06)	5.89 (4.96-6.81)	4.75 (3.67-6.04)	3.17 (1.91-2.96)	Missing
2000	7.57 (7.27-7.88)	6.5 (5.53-7.46)	4.67 (3.61-5.95)	3.3 (2.07-4.99)	Missing
2001	6.41 (6.14-6.69)	5.64 (4.75-6.52)	4.05 (3.1-5.19)	3.57 (2.37-5.17)	Missing
2002	5.55 (5.29-5.8)	5.45 (4.58-6.32)	3.32 (2.47-4.36)	3.12 (2.04-4.57)	Missing
2003	4.66 (4.43-4.9)	3.68 (2.97-4.39)	2.95 (2.18-3.89)	Missing	Missing
2004	3.77 (3.56-3.98)	3.47 (2.81-4.23)	2.8 (2.08-3.68)	2.59 (1.66-3.86)	Missing
2005	3.46 (3.26-3.66)	2.42 (1.89-3.06)	2.3 (1.66-3.11)	Missing	Missing
2006	2.84 (2.66-3.02)	2.13 (1.64-2.73)	1.76 (1.21-2.48)	Missing	Missing
2007	2.5 (2.33-2.66)	1.7 (1.27-2.23)	2.27 (1.66-3.03)	2.15 (1.38-3.21)	Missing
2008	2.31 (2.15-2.47)	2.09 (1.61-2.67)	2.32 (1.72-3.06)	Missing	Missing
2009	2.08 (1.93-2.23)	1.59 (1.18-2.09)	1.55 (1.09-2.14)	Missing	Missing
2010	2 (1.85-2.14)	1.76 (1.33-2.28)	1.93 (1.42-2.57)	1.58 (0.98-2.42)	Missing
2011	1.87 (1.73-2.01)	1.38 (1.01-1.83)	1.66 (1.21-2.23)	Missing	Missing
2012	1.87 (1.73-2.01)	1.79 (1.39-2.28)	1.65 (1.21-2.2)	Missing	Missing
2013	1.87 (1.73-2.01)	1.44 (1.08-1.88)	1.55 (1.14-2.06)	1.48 (0.95-2.2)	Missing
2014	1.79 (1.65-1.92)	1.49 (1.14-1.93)	2.38 (1.88-2.99)	Missing	Missing
2015	1.9 (1.76-2.03)	1.93 (1.53-2.41)	2.24 (1.77-2.8)	1.17 (0.75-1.74)	Missing
2016	2.74 (2.58-2.9)	1.99 (1.58-2.48)	3.13 (2.56-3.71)	1.6 (1.1-2.25)	Missing
2017	3.01 (2.84-3.18)	2.59 (2.11-3.07)	2.81 (2.27-3.36)	2.43 (1.82-3.18)	Missing
2018	3.28 (3.11-3.45)	2.44 (1.98-2.9)	3.47 (2.89-4.06)	1.58 (1.12-2.17)	Missing
2019	3.33 (3.16-3.5)	2.74 (2.26-3.22)	3.28 (2.73-3.84)	2.23 (1.68-2.91)	Missing
2020	4.02 (3.84-4.21)	3.21 (2.7-3.72)	5.06 (4.39-5.73)	2.59 (2.01-3.29)	Missing
2021	4.52 (4.31-4.72)	3.66 (3.1-4.21)	4.81 (4.16-5.47)	3.68 (2.95-4.53)	Missing
2022	4.15 (3.96-4.34)	2.94 (2.45-3.42)	4.32 (3.71-4.93)	2.79 (2.19-3.5)	Missing
2023	3.86 (3.67-4.04)	3.12 (2.61-3.62)	3.73 (3.17-4.29)	2.53 (1.97-3.2)	Missing

NH = non-Hispanic.

Supplemental Table 6 Angina Pectoris –related Crude Mortality Rates per 100,000, to 2000 U.S. population Stratified by Age Group in Adults in the United States, 1999 to 2023.

Year	Age 65-74 Years	75-84 Years	85+ Years
1999	2.54 (2.31-2.77)	7.26 (6.78-7.73)	29.51 (27.86-31.17)
2000	2.34 (2.12-2.56)	7.43 (6.95-7.92)	28.63 (27.02-30.25)
2001	2.24 (2.02-2.45)	6.32 (5.88-6.76)	23.47 (22.02-24.91)
2002	1.93 (1.73-2.13)	5.43 (5.02-5.83)	20.65 (19.3-21.99)
2003	1.51 (1.34-1.69)	4.41 (4.05-4.77)	17.62 (16.39-18.85)
2004	1.32 (1.16-1.49)	3.56 (3.24-3.89)	14.45 (13.35-15.56)
2005	1.07 (0.92-1.22)	3.17 (2.86-3.47)	13.27 (12.23-14.32)
2006	0.9 (0.77-1.04)	2.54 (2.26-2.81)	11.16 (10.22-12.1)
2007	0.83 (0.71-0.96)	2.2 (1.95-2.45)	10.18 (9.3-11.06)
2008	0.81 (0.69-0.93)	2.2 (1.95-2.46)	8.56 (7.77-9.36)
2009	0.72 (0.61-0.83)	2.13 (1.88-2.38)	7.4 (6.67-8.12)
2010	0.64 (0.53-0.75)	2 (1.76-2.24)	7.57 (6.84-8.3)
2011	0.68 (0.57-0.79)	1.8 (1.57-2.03)	6.48 (5.83-7.14)
2012	0.67 (0.56-0.77)	1.84 (1.61-2.07)	6.91 (6.24-7.58)
2013	0.65 (0.55-0.75)	1.82 (1.59-2.05)	6.84 (6.18-7.5)
2014	0.61 (0.52-0.71)	1.73 (1.51-1.95)	6.47 (5.84-7.11)
2015	0.77 (0.66-0.87)	1.9 (1.67-2.12)	6.68 (6.04-7.32)
2016	0.86 (0.75-0.97)	2.54 (2.28-2.81)	10.5 (9.71-11.3)
2017	1.03 (0.92-1.15)	2.92 (2.64-3.19)	11.16 (10.35-11.98)
2018	1.07 (0.96-1.19)	3.2 (2.91-3.48)	11.78 (10.95-12.61)
2019	1.14 (1.03-1.26)	3.15 (2.87-3.42)	12.6 (11.74-13.45)
2020	1.3 (1.17-1.42)	3.82 (3.52-4.12)	15 (14.07-15.93)
2021	1.49 (1.36-1.62)	4.33 (4.01-4.65)	16.86 (15.82-17.91)
2022	1.27 (1.15-1.39)	3.85 (3.56-4.14)	15.83 (14.86-16.8)
2023	1.18 (1.06-1.29)	3.51 (3.24-3.79)	15.25 (14.27-16.22)

Supplemental Table 7 Angina Pectoris–related Age-Adjusted Mortality Rates per 100,000, to 2000 U.S. population Stratified by Census Region in Older Adults in the United States, 1999 to 2023.

Census Region	Year	Age-Adjusted Rate
Northeast	1999	8.39 (7.73-9.05)
Northeast	2000	7.47 (6.86-8.09)
Northeast	2001	6.69 (6.11-7.27)
Northeast	2002	5.04 (4.54-5.55)
Northeast	2003	4.36 (3.89-4.82)
sNortheast	2004	3.82 (3.39-4.25)
Northeast	2005	3.4 (2.99-3.81)
Northeast	2006	2.49 (2.14-2.84)
Northeast	2007	2.29 (1.96-2.62)
Northeast	2008	1.76 (1.47-2.04)
Northeast	2009	1.67 (1.39-1.96)
Northeast	2010	1.62 (1.35-1.9)
Northeast	2011	1.35 (1.11-1.6)
Northeast	2012	1.27 (1.04-1.5)
Northeast	2013	1.35 (1.11-1.59)
Northeast	2014	1.09 (0.88-1.33)
Northeast	2015	1.23 (1-1.47)
Northeast	2016	1.78 (1.51-2.06)
Northeast	2017	1.99 (1.7-2.27)
Northeast	2018	2.16 (1.87-2.46)
Northeast	2019	2.24 (1.95-2.54)
Northeast	2020	2.95 (2.62-3.29)
Northeast	2021	3.03 (2.68-3.38)
Northeast	2022	2.44 (2.14-2.75)
Northeast	2023	2.38 (2.08-2.68)
Midwest	1999	7.8 (7.2-8.4)
Midwest	2000	7.52 (6.93-8.1)
Midwest	2001	6.58 (6.04-7.12)
Midwest	2002	5.72 (5.22-6.23)
Midwest	2003	4.2 (3.77-4.63)
Midwest	2004	3.71 (3.31-4.11)
Midwest	2005	3.13 (2.77-3.5)

Midwest	2006	2.56 (2.23-2.89)
Midwest	2007	2.48 (2.15-2.8)
Midwest	2008	1.96 (1.68-2.24)
Midwest	2009	1.62 (1.36-1.88)
Midwest	2010	1.72 (1.46-1.99)
Midwest	2011	1.46 (1.22-1.7)
Midwest	2012	1.51 (1.27-1.75)
Midwest	2013	1.38 (1.15-1.61)
Midwest	2014	1.41 (1.18-1.64)
Midwest	2015	1.48 (1.25-1.71)
Midwest	2016	2 (1.73-2.28)
Midwest	2017	2.14 (1.86-2.42)
Midwest	2018	2.4 (2.11-2.69)
Midwest	2019	2.57 (2.27-2.87)
Midwest	2020	2.97 (2.65-3.28)
Midwest	2021	3.42 (3.06-3.77)
Midwest	2022	2.91 (2.59-3.22)
Midwest	2023	2.87 (2.55-3.18)
South	1999	7.75 (7.25-8.26)
South	2000	8.27 (7.76-8.79)
South	2001	6.71 (6.25-7.18)
South	2002	5.8 (5.37-6.23)
South	2003	5.1 (4.71-5.5)
South	2004	3.77 (3.43-4.11)
South	2005	3.33 (3.01-3.64)
South	2006	2.8 (2.51-3.08)
South	2007	2.37 (2.11-2.62)
South	2008	2.45 (2.19-2.71)
South	2009	2.03 (1.79-2.26)
South	2010	1.83 (1.61-2.05)
South	2011	1.75 (1.54-1.97)
South	2012	1.98 (1.75-2.2)
South	2013	1.8 (1.6-2.01)
South	2014	1.82 (1.61-2.02)
South	2015	1.82 (1.62-2.02)
South	2016	2.5 (2.27-2.73)

South	2017	2.83 (2.59-3.08)
South	2018	3.19 (2.93-3.44)
South	2019	3.48 (3.21-3.74)
South	2020	3.89 (3.61-4.16)
South	2021	4.46 (4.16-4.77)
South	2022	4.37 (4.08-4.66)
South	2023	4.13 (3.85-4.42)
West	1999	5.77 (5.19-6.35)
West	2000	5.48 (4.92-6.03)
West	2001	4.77 (4.26-5.29)
West	2002	4.97 (4.45-5.48)
West	2003	4.12 (3.65-4.58)
West	2004	3.49 (3.07-3.92)
West	2005	3.31 (2.9-3.72)
West	2006	3.11 (2.72-3.51)
West	2007	2.81 (2.44-3.17)
West	2008	2.77 (2.41-3.13)
West	2009	2.92 (2.56-3.29)
West	2010	2.73 (2.38-3.07)
West	2011	2.62 (2.28-2.96)
West	2012	2.54 (2.21-2.86)
West	2013	2.71 (2.38-3.04)
West	2014	2.54 (2.22-2.86)
West	2015	2.91 (2.58-3.24)
West	2016	4.21 (3.81-4.6)
West	2017	4.76 (4.34-5.17)
West	2018	4.59 (4.19-4.99)
West	2019	4.44 (4.05-4.83)
West	2020	5.52 (5.09-5.95)
West	2021	6.33 (5.85-6.8)
West	2022	5.69 (5.25-6.12)
West	2023	4.99 (4.58-5.4)

Supplemental Table 8 Angina Pectoris–related Age-Adjusted Mortality Rates per 100,000, to 2000 U.S. population Stratified by Urban-Rural Classification in Older Adults in the United States, 1999 to 2020.

Year	Metropolitan	Non-metropolitan
1999	6.98 (6.6-7.29)	9.82 (9.08-10.57)
2000	6.75 (6.45-7.06)	9.98 (9.22-10.73)
2001	5.49 (5.22-5.76)	9.48 (8.75-10.21)
2002	4.87 (4.62-5.13)	8.04 (7.36-8.71)
2003	4.03 (3.8-4.26)	6.64 (6.03-7.25)
2004	3.31 (3.1-3.52)	5.65 (5.09-6.2)
2005	2.99 (2.79-3.18)	4.75 (4.24-5.26)
2006	2.49 (2.31-2.66)	3.8 (3.35-4.26)
2007	2.32 (2.15-2.48)	3.1 (2.7-3.51)
2008	2.04 (1.88-2.19)	3.25 (2.83-3.67)
2009	1.85 (1.7-1.99)	2.78 (2.4-3.16)
2010	1.83 (1.69-1.98)	2.65 (2.28-3.02)
2011	1.69 (1.56-1.83)	2.3 (1.96-2.64)
2012	1.71 (1.58-1.85)	2.5 (2.14-2.85)
2013	1.77 (1.64-1.91)	2.16 (1.84-2.49)
2014	1.68 (1.54-1.81)	2.15 (1.83-2.47)
2015	1.83 (1.69-1.96)	2.23 (1.91-2.56)
2016	2.65 (2.49-2.81)	2.69 (2.33-3.05)
2017	2.81 (2.64-2.97)	3.65 (3.24-4.06)
2018	3.05 (2.88-3.21)	3.49 (3.09-3.88)
2019	3.21 (3.04-3.38)	3.66 (3.26-4.06)
2020	3.71 (3.53-3.89)	4.58 (4.13-5.03)